# UnitedHealthcare Medicare Advantage prescription drug plans: Opioid overutilization prevention

Quick reference guide for pharmacy

Please use this quick reference guide to learn more about how to work with our UnitedHealthcare® Medicare Advantage prescription drug plan pharmacy programs. These programs can help your patients get the opioid treatments they need in safe and effective ways.

We use opioid safety edits at point-of-sale (POS) to prompt prescribers and pharmacists to conduct additional safety reviews to determine if the member's opioid use is appropriate and medically necessary. Our goal is to implement these edits in a manner that helps minimize any additional burden on prescribers, pharmacists and our members.

For more information about resolving rejected prescription claims that have opioid safety edits or plan dispensing limits, please access the **Optum Rx Pharmacy Provider Manual** at **professionals.optumrx.com**.



## **Pharmacy-based programs**

- **1** Opioid Dispensing Limit
- 2 Concurrent Drug Utilization Review (cDUR) (opioid safety alerts)
- 3 Retrospective Drug Utilization Review: Drug Management Program (DMP)



### **Opioid Dispensing Limit**

This program helps support safe and appropriate opioid use while limiting excess quantities.

Plan benefit	Description and messaging	Pharmacy action
Opioid Dispensing Limit	Limits opioid dispensing by mail and at retail locations to a 1-month supply per prescription.  POS message: Maximum Days' Supply "displays plan limit"	Pharmacy cannot override the dispensing limit.  Reprocess the prescription claim for a 1-month supply (day supply in reject messaging).
	Reject codes: 76: Plan Limitations Exceeded 19: M/I Days Supply	





# **Concurrent Drug Utilization Review (cDUR) (opioid safety alerts)**

The cDUR program uses the pharmacy claims processing system to screen all prescriptions at POS. It screens for inappropriate drug prescriptions and utilization, as well as potentially dangerous medical implications or medication interactions. We communicate with the dispensing pharmacy at POS through claims messaging.

Opioid safety alert	Description and messaging	Pharmacy action
7-day supply limit for opioid naïve patients	We'll limit members to a supply of 7 days or less if they haven't filled an opioid prescription within 120 days.  POS message: 7 day max opioid naïve  Reject codes: 88+925	If the member has historical opioid use: Resolve at POS with pharmacist intervention.  We allow the following resolution code combinations:  MX, M0, 4J Prescriber consulted; patient isn't opioid naïve  MX, R0, 4J Pharmacist consulted; patient isn't opioid naïve  MX, MR, 4J Medication review; patient isn't opioid naïve  If the patient doesn't have historical
		<ul> <li>opioid use:</li> <li>1. The pharmacy resubmits the claim for a 7-day supply only.</li> <li>2. The pharmacy may override the rejection when an exemption exists. Use standard NCPDP Prospective Payment System (PPS) codes to resolve at POS.</li> <li>3. We require prior authorization for quantities of more than 7-day supplies that don't meet the above criteria.</li> </ul>
Opioid care coordination (OCC) safety alert	Limits cumulative morphine milligram equivalent (MME)* dosage per day for all opioid prescriptions when the threshold exceeds 90 MME and the patient receives opioid prescriptions from more than 1 prescriber.  POS message: MME ### Exceeded; Total MME XXXmg, OCC: MD Consult use: HD-M0-1G, Exempt: use PPS codes; RxHelp 800-788-7871, Consider Naloxone for safety  Reject code(s): 88+922	If the pharmacist recently consulted with the prescriber and has up-to-date clinical information (e.g., Prescription Drug Monitoring Program (PDMP) system or other records), we don't expect additional consultation with the prescriber.  Prescriber consultation: Enter HD-M0-1G.  Pharmacist may override the rejection when an exemption exists. Exemptions may depend on pharmacy drug claim history, knowledge of the patient's diagnosis and prescriber specialty. Use standard NCPDP PPS codes to resolve at POS.  Important note: Pharmacies should only use the override code 1G after completing and documenting the care coordination activities with prescribers. We have the option to audit pharmacies' documentation.



Opioid safety alert	Description and messaging	Pharmacy action
Cumulative MME opioid safety alert	Limits cumulative MME dosage per day for opioid prescriptions when the threshold exceeds 200 MME and the patient receives opioid prescriptions from more than 1 prescriber.	The pharmacy may override the reject when an exemption exists. Use standard NCPDP PPS codes to resolve at POS.  We require prior authorization in all other instances.
	POS message: MME ### Exceeded; Total MME XXXmg, OPIOID MME PA REQ'D, 800-711-4555, Exempt: use PPS codes, Consider Naloxone for safety	
	Reject codes: 88+922+G4	
Opioids and buprenorphine (used in medication-assisted treatment (MAT))	Alert for concurrent use of opioids and buprenorphine.	Pharmacist conducts an additional safety review to determine if the patient's concurrent opioid use is safe and clinically appropriate. Use standard NCPDP PPS codes to resolve at POS.
	The pharmacist will conduct additional safety reviews to determine if the patient's concurrent use is safe and clinically appropriate. The pharmacist may contact the prescriber.	
	POS message: Opioid+Buprenorphine Hx, MD Cnslt: use DD-M0-1G; Exempt: use PPS code, RxHelp 800-788-7871	
	Reject code: 88	
Opioids and benzodiazepines	Alert for concurrent use of opioids and benzodiazepines.	Pharmacist conducts an additional safety review to determine if the patient's concurrent opioid use is safe and clinically appropriate. Use standard NCPDP PPS codes to resolve at POS.
	The pharmacist will conduct additional safety reviews to determine if the patient's opioid use is safe and clinically appropriate. The pharmacist may contact the prescriber.	
	POS message: Benzo+OpioidHx; Consider Naloxone MD Cnslt: use DD-M0-1G; Exempt: use PPS code; RxHelp 800-788-7871	
	Reject code: 88	
Duplicative long-acting opioid (LAO) therapy	Alert for concurrent use of multiple LAOs.	Pharmacist conducts additional safety review to determine if the patient's opioid use is safe and clinically appropriate. Use standard NCPDP PPS codes to resolve at POS.
	Pharmacist conducts an additional safety review to determine if the patient's opioid use is safe and clinically appropriate. The pharmacist may contact the prescriber.	
	POS message: LA Opioid Dup Therapy, MD Cnslt: use TD-M0-1G; Exmpt:use PPS code; RxHelp 800-788-7871	
	Reject code: 88	





## Retrospective Drug Utilization Review: Drug Management Program (DMP)

The Drug Management Program helps address overutilization of frequently abused drugs while maintaining member access to the medications as medically necessary. Each month, we identify the members who meet Medicare Part D Overutilization Monitoring System (OMS) criteria, such as members who are at-risk of obtaining opioids from multiple prescribers or pharmacies. We also review the prescriptions of members with histories of opioid-related overdose.

Plan benefit	Description and messaging	Pharmacy action
Drug Management Program (DMP)	Reject codes:  828 - Beneficiary Case Management Restriction  980 - Patient Locked into Specific Pharmacy(s)  979 - Patient Locked into Specific Prescriber(s)  POS message:  Fax Written Request 877-275-6030  *Multiple reject codes are possible if multiple restrictions are in place (e.g., pharmacy and prescriber lock in 979 + 980).	Pharmacy cannot override coverage limitations or restrictions resulting from our DMP review.  To request changes to imposed coverage limitations or restrictions, please fax us at 877-275-6030. Or, the member may contact Optum Case Management at 855-218-3456 for ongoing case management review.



#### Reminders

#### Drug utilization review (DUR) intervention exemptions

- Residents of long-term care facilities
- · Patients receiving hospice care
- · Patients receiving palliative or end-of-life care
- Patients receiving treatment for active cancer-related pain or sickle cell disease

Please note that opioid safety edits don't impact patients' access to MAT.

#### **Clinical considerations**

Consider offering naloxone when factors that increase risk for opioid overdose are present, such as:

- · History of overdose
- · History of substance use disorder
- Higher opioid dosages (more than 50 MME/day)
- Concurrent benzodiazepine use

#### **Resolving POS rejects**

We encourage pharmacies to resolve safety edits expeditiously and avoid disruption of therapy. Please only submit prior authorization when it's necessary to resolve a POS reject. In this case, you can submit a prior authorization request using one of the following options:

- Visit professionals.optumrx.com
- Call Optum Rx prior authorization at 800-711-4555, available 5 a.m.–10 p.m. PT, Monday–Friday and 6 a.m.–3 p.m. PT, Saturday



## **Questions?**

If you have questions, please contact the Optum Rx Pharmacy Help Desk, available 24 hours a day, 7 days a week at:

- 877-889-6510 (Medicare Advantage Prescription Drug Plan (MA-PD))
- 877-889-6481 (Medicare Prescription Drug Plan (PDP))

