

Brineura® (Cerliponase Alfa)

Policy Number: CS2023D0065M
Effective Date: August 1, 2023

[Instructions for Use](#)

Table of Contents	Page
Application	1
Coverage Rationale	1
Applicable Codes	2
Background	2
Clinical Evidence	3
U.S. Food and Drug Administration	4
References	4
Policy History/Revision Information	4
Instructions for Use	4

Commercial Policy
<ul style="list-style-type: none"> Brineura® (Cerliponase Alfa)

Application

This Medical Benefit Drug Policy does not apply to the states listed below; refer to the state-specific policy/guideline, if noted:

State	Policy/Guideline
Florida	Refer to the state’s Medicaid clinical policy
Indiana	Brineura® (Cerliponase Alfa) (for Indiana Only)
Kansas	Refer to the state’s Medicaid clinical policy
Louisiana	Brineura® (Cerliponase Alfa) (for Louisiana Only)
North Carolina	None
Ohio	Brineura® (Cerliponase Alfa) (for Ohio Only)
Pennsylvania	Brineura® (Cerliponase Alfa) (for Pennsylvania Only)
Texas	Refer to drug-specific criteria found within the Texas Medicaid Provider Procedures Manual

Coverage Rationale

Brineura (cerliponase alfa) is proven and medically necessary for slowing the loss of ambulation in symptomatic pediatric patients with late infantile neuronal ceroid lipofuscinosis (LINCL) type 2 (CLN2), also known as tripeptidyl peptidase 1 (TPP1) deficiency, when all of the following criteria are met:^{1-6,10-15}

- For **initial therapy**, all of the following:
 - **One** of the following:
 - Diagnosis of late infantile neuronal ceroid lipofuscinosis type 2 (CLN2) by a neurologist with expertise in the diagnosis of CLN2
 - Diagnosis of late infantile neuronal ceroid lipofuscinosis type 2 (CLN2) by a physician in consultation with a neurologist with expertise in the diagnosis of CLN2
 - and**
 - Patient is age 3 years or older; **and**
 - **All** of the following scores on the Clinical Scoring System for LINCL:⁴

- Combined score of 3 to 6 in the motor and language domains
- Score of at least 1 in the motor domain
- Score of at least 1 in the language domain

and

- **One** of the following:

- Brineura is prescribed by a neurologist with expertise in the treatment of CLN2
- Brineura is prescribed by a physician in consultation with a neurologist with expertise in the treatment of CLN2

and

- Brineura is to be administered intraventricularly by, or under the direction of, healthcare professionals experienced in performing intraventricular infusions via an intracerebroventricular catheter; **and**
- Dosing is in accordance with the U.S. Food and Drug Administration (FDA) approved labeling; **and**
- Initial authorization will be for no more than 6 months

- For **continuation of therapy**, **all** of the following:

- **One** of the following:

- Diagnosis of late infantile neuronal ceroid lipofuscinosis type 2 (CLN2) by a neurologist with expertise in the diagnosis of CLN2
- Diagnosis of late infantile neuronal ceroid lipofuscinosis type 2 (CLN2) by a physician in consultation with a neurologist with expertise in the diagnosis of CLN2

and

- Patient is age 3 years or older; **and**
- Patient has a score of 1 or higher in the motor domain of the Clinical Scoring System for LINCL;^{4,7} ; **and**
- **One** of the following:

- Brineura is prescribed by a neurologist with expertise in the treatment of CLN2
- Brineura is prescribed by a physician in consultation with a neurologist with expertise in the treatment of CLN2

and

- Brineura is to be administered intraventricularly by, or under the direction of, healthcare professionals experienced in performing intraventricular infusions via an intracerebroventricular catheter; **and**
- Dosing is in accordance with the U.S. Food and Drug Administration (FDA) approved labeling; **and**
- Reauthorization will be for no more than 6 months

Brineura (cerliponase alfa) is unproven and not medically necessary for other forms of neuronal ceroid lipofuscinosis.

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state, or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

HCPCS Code	Description
J0567	Injection, cerliponase alfa, 1 mg

Diagnosis Code	Description
E75.4	Neuronal ceroid lipofuscinosis

Background

Neuronal ceroid lipofuscinosis type 2 (CLN2) is a neurodegenerative lysosomal storage disorder caused by deficient activity of the enzyme tripeptidyl peptidase-1 (TPP1).¹ Deficiency in TPP1 activity results in the accumulation of lysosomal storage materials normally metabolized by this enzyme in the central nervous system (CNS), leading to progressive decline in motor function. CLN2 is autosomal recessive and pediatric-onset, and is characterized by seizures, language delay, movement disorders, motor deterioration, dementia, blindness, and early death.^{2,3} A Clinical Scoring System for late infantile neuronal

ceroid lipofuscinoses has been developed as a method for quantitative description of clinical courses over time.⁴ Within CLN2, two forms of disease evolution exist; classical CLN2 is where symptoms start earlier, between the ages of 3 and 5 years and the symptoms evolve faster.⁷ Non-classical CLN2 has a much slower disease evolution and symptoms appear as behavioral disorders, movement disorders and ataxia rather than seizures and blindness.

Cerliponase alfa (rhTTP1), a proenzyme, is taken up by target cells in the CNS and is translocated to the lysosomes through the Cation Independent Mannose-6-Phosphate Receptor (CI-MPR, also known as M6P/IGF2 receptor). Cerliponase alfa is activated in the lysosome and the activated proteolytic form of rhTTP1 cleaves tripeptides from the N-terminus of proteins.

Clinical Evidence

Proven

Ceroid Lipofuscinosis Type 2 (CLN2)/Tripeptidyl Peptidase 1 (TPP1) Deficiency

Cerliponase alfa is indicated to slow the loss of ambulation in symptomatic pediatric patients 3 years of age and older with late infantile neuronal ceroid lipofuscinosis type 2 (CLN2), also known as tripeptidyl peptidase 1 (TPP1) deficiency.¹

In a multicenter, open-label study, Schulz A. et al. evaluated the effect of intraventricular infusion of cerliponase alfa every 2 weeks in pediatric patients with CLN2.⁶ The primary outcome compared the duration until a 2-point decline in the score on the motor and language domains of the CLN2 Clinical Rating Scale in study patients to the rate of decline in 42 historical controls. In addition, the rate of decline in the motor-language score was compared between the two groups. Of the 24 patients enrolled, 23 constituted the efficacy population. The median time until a 2-point decline in the motor-language score was not reached for treated patients and was 345 days for historical controls. The mean (\pm SD) unadjusted rate of decline in the motor-language score per 48-week period was 0.27 ± 0.35 points in treated patients and 2.12 ± 0.98 points in 42 historical controls (mean difference, 1.85; $p < 0.001$). Common adverse events included convulsions, pyrexia, vomiting, hypersensitivity reactions, and failure of the intraventricular device. Infections developed in the intraventricular device for administration in two patients, required antibiotic treatment and device replacement. The authors conclude that intraventricular infusion of cerliponase alfa in patients with CLN2 disease resulted in less decline in motor and language function than that in historical controls.

Clinical evidence for the safety and efficacy of cerliponase alfa for the treatment of late infantile neuronal ceroid lipofuscinosis type 2 (CLN2) was demonstrated in a prospective Phase 1/2 Open-Label Dose-Escalation Study and Extension. The objective of the study was to evaluate the safety and tolerability of cerliponase alfa administered to patients with CLN2 disease by intraventricular administration. There were five study centers involved. Patients were treated with intraventricular infusion of cerliponase alfa with doses ranging from 30 to 300 mg every 14 days in the dose escalation study and were maintained at 300 mg every 14 days in the extension study. The primary endpoint was response rate, defined as the absence of an unreversed two-point decline or score of zero in the CLN2 score at 48 weeks. 24 patients were enrolled, with 23 patients completing the study. By motor/language CLN2 scores measured from baseline, 87% (20/23) of treated patients responded to treatment, defined as an absence of an unreversed two-point decline or score of zero by Week 48, compared to an expected response rate of 50% ($p = 0.0002$). Sixty-five percent of treated patients experienced no progression in their CLN2 score. Of all points lost, approximately 80% occurred within four months of treatment initiation. The proportion of patients with a response to treatment was 87% at Week 48 and 63% at Week 96.⁵

Twenty-one international experts from seven different specialties developed guidelines on the diagnosis, clinical assessments, treatment, and management for CLN2 disease patients that were published in 2021.⁷ From a consensus statement in these guidelines, it is stated that initiation of long-term ERT with cerliponase alfa at 300 mg (or age-appropriate) dose every other week through intraventricular infusion is suggested in non-classical TPP1 deficiency patients after confirmed diagnosis and agreement between parents and provider, as long as no contraindications to therapy exist. Furthermore, initiation of long-term ERT with cerliponase alfa at 300 mg (or age-appropriate) dose every other week through intraventricular infusion is recommended in classical CLN2 patients with the potential to benefit from this therapy.

U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Brineura (cerliponase alfa) is a hydrolytic lysosomal N-terminal tripeptidyl peptidase indicated to slow the loss of ambulation in symptomatic pediatric patients 3 years of age and older with late infantile neuronal ceroid lipofuscinosis type 2 (CLN2), also known as tripeptidyl peptidase 1 (TPP1) deficiency.¹

References

1. Brineura [package insert]. Novato, CA: BioMarin Pharmaceutical Inc.; March 2020.
2. Williams RE, Adams HR, Blohm M, Cohen-Pfeffer JL, de Los Reyes E, Denecke J, et al. Management Strategies for CLN2 Disease. *Pediatr Neurol*. 2017 Apr;69:102-112.
3. <http://www.cln2connection.com/overview/cln2-disease>. Accessed June 6, 2023.
4. Steinfeld R, Heim P, von Gregory H, et al. Late infantile neuronal ceroid lipofuscinosis: quantitative description of the clinical course in patients with CLN2 mutations. *Am J Med Genet*. 2002;112:347-354.
5. AMCP Dossier for Brineura® (cerliponase alfa), BioMarin Pharmaceutical, May 2017.
6. Schulz A, et al. Study of Intraventricular Cerliponase Alfa for CLN2 Disease. *N Engl J Med*. 2018 Apr 24.
7. Mole SE, Schulz A, Badoe E, et al. Guidelines on the diagnosis, clinical assessments, treatment and management for CLN2 disease patients. *Orphanet J Rare Dis*. 2021;16(1):185. Published 2021 Apr 21. doi:10.1186/s13023-021-01813-5.

Policy History/Revision Information

Date	Summary of Changes
08/01/2023	<ul style="list-style-type: none">• Routine review; no change to coverage guidelines• Archived previous policy version CS2023D0065L

Instructions for Use

This Medical Benefit Drug Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state or contractual requirements for benefit plan coverage govern. Before using this policy, please check the federal, state or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Benefit Drug Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the InterQual® criteria, to assist us in administering health benefits. The UnitedHealthcare Medical Benefit Drug Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.